

Extreme emission line outflows in the GPS source 4C 12.50 (PKS 1345+12)

J. Holt,¹ C. N. Tadhunter,¹ and R. Morganti²

¹ The University of Sheffield, Hicks Building, Hounsfield Road, Sheffield, S3 7RH, UK.
j.holt@sheffield.ac.uk, c.tadhunter@sheffield.ac.uk

² ASTRON, PO Box 2, 7990 AA Dwingeloo, The Netherlands.
morganti@astron.nl

Abstract

We present high resolution spectra ($0.7 \text{ \AA}/\text{pix}$) of the GPS source 4C 12.50 with large spectral coverage ($\sim 4500 \text{ \AA}$) taken with the 4.2m William Herschel Telescope, La Palma. The slit was aligned along PA 160 to include the nucleus and emission line region to the NW. An asymmetric halo extending 20 kpc NW and 12 kpc SE from the nucleus is clearly seen. At the position of the nucleus we observe unusually broad forbidden emission line components (broadest component: $\text{FWHM} \sim 2000 \text{ km s}^{-1}$), blue shifted by up to 2000 km s^{-1} with respect to the halo of the galaxy and H I absorption. We interpret this as material in outflow. We measure $E(B-V) = 1.44$ for the broadest, most kinematically disturbed component, corresponding to an actual $\text{H}\beta$ flux 130 times brighter than that measured. We calculate an upper limit for the mass of the line emitting gas of order $10^6 M_{\odot}$ for both the intermediate and broad components. Our results are consistent with 4C 12.50 being a young radio source.

Keywords: ISM: jets and outflows - ISM: kinematics and dynamics - galaxies: active - galaxies: ISM - galaxies: kinematics and dynamics - galaxies: individual (4C 12.50, PKS 1345+12)

1 Introduction

The narrow line region ($\text{NLR} < 5 \text{ kpc}$) is an important probe of the ISM close to the central engine in radio galaxies (Tadhunter *et al.*, 2001).

Powerful *extended* radio galaxies have kinematics generally consistent with gravitational motions in the bulges of the host galaxies (e.g. Heckman *et al.*, 1984). In contrast, GPS radio sources ($< 1 \text{ kpc}$ in size) show evidence for kinematic disturbance (Gelderman & Whittle, 1994). Highly complex broad emission lines are observed but the nature of the emission line kinematics is still not understood. Asymmetries are believed to be the spectral signatures of systematic flows, but their direction (outflow or inflow) is still under dispute due to the lack of understanding of the dust distribution in the NLR, and lack of accurate host galaxy (rest frame) redshifts.

Compact radio sources are thought to be either young radio sources (e.g. Fanti *et al.*, 1995) or intrinsically compact objects confined by an unusually dense ISM (the ‘frustration theory’, e.g. van Breugel, 1984). High reddening is expected in both scenarios - the nucleus is likely to be enshrouded by large amounts of gas and dust left over from the event(s) which triggered the activity (e.g. a merger, Heckman *et al.*, 1986). With evolution, material along the radio axis will be dissipated by jet-cloud interactions and/or quasar-induced winds (Tadhunter *et al.*, 2001). Eventually cavities will be hollowed out on either side of the nucleus, as is seen in the powerful extended radio source Cygnus A (Tadhunter *et al.*, 1999).




Figure 1: High contrast HST WFPC2 image in [O III] from Axon *et al.* (2000) with the VLA radio map (5 GHz) from Stanghellini *et al.* (1997). For a higher resolution map, refer to the original publication.

2 Previous observations

4C 12.50 has a double nucleus and shows clear evidence for distorted morphology in the ionised gas: high resolution HST images reveal an arc-like structure 1.2 arcsec (2.8 kpc) N of the nucleus and fainter diffuse emission 2 arcsec (4.7 kpc) N of the nucleus (see figure 1). The radio source has distorted, triple morphology aligned along PA 160 with emission confined to a region < 0.1 arcsec (~ 240 pc) in size (see figure 1). A bright knot towards the northern limit of the jet is tentatively identified as the core (Stanghellini *et al.*, 1997). The jet extends 0.04 arcsec (95 pc) to the SE before bending and expanding into a diffuse lobe. Weak emission extends up to 0.03 arcsec (70 pc) NW of the core. ($H_0 = 75 \text{ km s}^{-1}$, $q_0 = 0.0$).

3 Results

These *preliminary* results are based on new, high resolution spectroscopic observations ($\sim 0.7 \text{ \AA/pix}$) with large spectral coverage ($\sim 4500 \text{ \AA}$) taken in 2001 May with the 4.2m William Herschel Telescope (WHT) on La Palma. For all observations, the slit width was 1.3 arcsec and the seeing was ~ 1.1 arcsec (FWHM) or better. For a complete analysis and discussion of PKS 1345+12, see Holt *et al.* (2002, in preparation).

3.1 Modelling the emission lines

The emission lines were modelled using the Starlink package DIPSO constraining, for example, the intensity ratios and shifts between doublet lines, where possible, in accordance with atomic physics. In order to model the unusually broad [O III] $\lambda\lambda 4959, 5007$ emission lines in the nucleus it is essential to use three Gaussians (see figure 2). These three components represent line emission from three distinct redshift systems:




Figure 2: Model for $[\text{O III}]\lambda\lambda 4959,5007$ (bold line) comprising 3 components for each line: narrow (dot-dashed line), intermediate (solid line) and broad (dotted line).

- i)* A narrow component ($\text{FWHM} \sim 360 \text{ km s}^{-1}$) assumed to emanate from the ambient, quiescent ISM. Its velocity is consistent with the deep HI 21cm absorption (Mirabel, 1989; see also Morganti *et al.*, these proceedings), and the extended $[\text{O II}]$ halo.
- ii)* An intermediate component ($\text{FWHM} \sim 1260 \text{ km s}^{-1}$) blue-shifted by $\sim 400 \text{ km s}^{-1}$ relative to the narrow component.
- iii)* A broad component ($\text{FWHM} \sim 1960 \text{ km s}^{-1}$) blue-shifted by $\sim 1980 \text{ km s}^{-1}$ relative to the narrow component.

The velocity shifts in the $[\text{O III}]$ lines clearly indicate that they originate from three distinct regions along the line-of-sight. The narrowest component of $[\text{O III}]$ is consistent with the spatially extended $[\text{O II}]$ emission and HI absorption and represents the galaxy rest frame. Hence, the intermediate and broad components of $[\text{O III}]$ trace material blue-shifted with respect to the galaxy system, suggesting that the ISM emitting these components is in outflow (see figure 3).

3.2 Evidence for high reddening in the nucleus

We measure $E(B-V)$ values, calculated from the $\text{H}\alpha/\text{H}\beta$ ratio, of 0.06, 0.42 and 1.44 for the narrow, intermediate and broad components respectively. The broad component is highly reddened and corrections for reddening yield $\text{H}\beta$ fluxes a factor of 130 brighter than those measured. This reddening is consistent with $\text{Pa}\alpha/\text{H}\alpha$ ratios estimated using $\text{Pa}\alpha$ measurements from Veilleux *et al.* (1997). Note, the highly broadened $\text{Pa}\alpha$ component detected by Veilleux *et al.* is not associated with a classical Broad Line Region (BLR) as it is also observed in the forbidden lines and therefore represents a broadened NLR component.




Figure 3: Comparison of the three components of [O III] (open triangles) with the spatially extended [O II] $\lambda\lambda 3726, 3729$ emission (open diamonds) and the deep HI 21cm absorption from Mirabel *et al.* (1989) (filled circle) provides a good estimate of the galaxy rest frame. Two of the [O III] components trace material in outflow.

3.3 Estimating the gas mass

Emission line luminosities are related to the mass of line emitting gas by:

$$M_{gas} = m_p \frac{L(H\beta)}{N_e \alpha_{H\beta}^{eff} h \nu_{H\beta}}$$

where N_e is the electron density (cm^{-3}); m_p is the mass of the proton (kg); $L(H\beta)$ is the luminosity of the $H\beta$ line: 0.44 ± 0.04 , 5.50 ± 0.20 and 26.59 ± 0.59 ($10^{40} \text{ erg s}^{-1}$) for the narrow, intermediate and broad components respectively; $\alpha_{H\beta}^{eff}$ is the effective recombination coefficient for $H\beta$ ($\text{cm}^3 \text{ s}^{-1}$; see Osterbrock 1989) and $h\nu_{H\beta}$ is the energy of the $H\beta$ photons (erg).

The density diagnostic, the [S II] $\lambda\lambda 6716, 6730$ doublet, can only be modelled with very high densities for the intermediate and broad components. We estimate lower limits for the density of the intermediate and broad components of $\sim 4000 \text{ cm}^{-3}$. These lead to an upper limit for the total mass of the line emitting gas of $9.17 \times 10^5 M_{\odot}$.

4 Conclusions

Our data show:

- complex broad emission lines;
- blue shifts up to $\sim 2000 \text{ km s}^{-1}$ with respect to the galaxy halo and HI absorption;
- large reddening (broadest component $E(B-V) \sim 1.44$);
- high densities and high ionisation for the intermediate and broad components.

For the blue shifted components we calculate:

- a lower limit for the density;
- an upper limit for the gas mass of order $10^6 M_{\odot}$.




Figure 4: Model proposed for the compact flat spectrum radio source PKS 1549-79 by Tadhunter *et al.* (2001).

Our results are consistent with the model proposed for the compact flat spectrum radio source PKS 1549-79 by Tadhunter *et al.* (2001) in which the nucleus is cocooned in gas and dust in the early stages and drives outflows which will eventually hollow out the ionisation cones on either side of the nucleus (see figure 4).

Our estimates for the mass of emitting gas of order $10^6 M_{\odot}$ are also consistent with the radio source being young. Using the equations presented by O’Dea (1998) from de Young (1993), a smooth medium with the density we measure would require gas masses of order $10^8 M_{\odot}$ to $10^{10} M_{\odot}$ to frustrate the radio source - much larger than we estimate for the blue shifted components.

Acknowledgements

JH acknowledges a PPARC PhD studentship. The William Herschel Telescope is operated on the island of La Palma by the Isaac Newton Group in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias.

References

- Axon D. J., Capetti A., Fanti R., Morganti R., Robinson A. & Spencer R., 2000, *AJ*, 120, 2284
- de Young D. S., 1993, *ApJ*, 402, 95
- Fanti C., Fanti R., Schilizzi R. T., Spencer R. E. & Stanghellini C., 1995, *A&A*, 302, 317
- Gelderman R. & Whittle M., 1994, *ApJS*, 91, 491
- Heckman T. M., Miley G. K. & Green R. F., 1984, *ApJ*, 281, 525
- Heckman T. M., Smith E. P., Baum S. A., van Breugel W. J. M., Miley G. K., Illingworth G. D., Bothun G. D. & Balick B., 1986, *ApJ*, 311, 526
- Mirabel I. F. et al, 1989, *ApJ*, 340, L13
- O’Dea, C. P., 1998, *PASP*, 110, 493
- Osterbrock D. E., 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*, University Science Books, p. 80
- Stanghellini C., O’Dea C. P., Baum S. A., Dallacasa D., Fanti R. & Fanti C., 1997, *A&A*, 325, 943
- Tadhunter C. N., Packham C., Axon D. J., Robinson A., Hough J., Young S., Sparks W., 1999, *ApJ*, 512, L91
- Tadhunter C. N., Wills K., Morganti R., Oosterloo T. & Dickson R., 2001, *MNRAS*, 327, 227
- van Breugel W. J. M., 1984, *IAU symp.*, 110, p.59
- Veilleux S., Sanders D. B. & Kim D.-C., 1997, *ApJ*, 484, 92